



WHAT IS CLAIMED IS:

1. A process for preparing (R)-hydroxy ester II5 from alpha-keto ester I

comprising adding the alpha-keto ester I to a mixture comprising the ketoreductase enzyme and non-ketoreductase enzyme components, wherein

10 R¹ is C₁₋₄ alkyl; andR² is selected from the group consisting ofC₁₋₈ alkyl,C₁₋₈ alkyl, substituted with C₃₋₇ cycloalkyl,

aryl, and

15 a 5- to 7-membered saturated or unsaturated heterocyclic ring,

wherein the ketoreductase enzyme has a molecular weight between 36000 and 38000, and wherein the ketoreductase enzyme has an N-terminal amino acid sequence selected from the group of sequences consisting of

20

Ala-Ile-Pro-Asp-Asn-Ala-Val-Leu-Glu-Gly-Ser-Leu-Val-Lys-Val-Thr-Gly-Ala-Asn-Gly (SEQ. ID NO. 1),

25 Met-Ala-Ile-Pro-Asp-Asn-Ala-Val-Leu-Glu-Gly-Ser-Leu-Val-Lys-Val-Thr-Gly-Ala-Asn-Gly (SEQ. ID NO. 4),

and

Met-Ala-Lys-Ile-Asp-Asn-Ala-Val-Leu-Pro-Glu-Gly-Ser-Leu-Val-Leu-Val-Thr-Gly-Ala-Asn-Gly (SEQ ID NO. 2).

5

2. A process of Claim 1, wherein the N-terminal amino acid sequence is selected from the group consisting of

10 Ala-Ile-Pro-Asp-Asn-Ala-Val-Leu-Glu-Gly-Ser-Leu-Val-Lys-Val-Thr-Gly-Ala-Asn-Gly, and

Met-Ala-Ile-Pro-Asp-Asn-Ala-Val-Leu-Glu-Gly-Ser-Leu-Val-Lys-Val-Thr-Gly-Ala-Asn-Gly.

15

3. A process of Claim 2, wherein the non-ketoreductase enzyme components comprise NADP, a cofactor recycling system comprising a hydride source and a catalyst, and a buffer suitable for maintaining a pH of between about 5 and about 10.

20

4. A process of Claim 3, wherein temperature is between about 25 and about 40°C.

25 5. A process of Claim 4, wherein the hydride source is glucose and the catalyst is glucose dehydrogenase.

6. A process of Claim 5, wherein the mixture comprises an amount of ketoreductase between about 0.1 and about 10 g/L, an amount of NADP between about 0.1 and about 10 g/L.

30

7. A process of Claims 1, wherein R² is selected from the group consisting of

